

A
S, NH or carbonyl, for example $C(O)R^1$, $CH_2C(O)R^1$, $NHCH_2R^1$, $NHC(O)R^1$, CH_2OR^1 , $CH_2C(O)NHR^1$, and the like.

REMARKS

Attached hereto is a marked-up version of the changes made to the specification and claims by the current amendment. The attached page is captioned "VERSION WITH MARKINGS TO SHOW CHANGES MADE." The Office is invited to contact the undersigned should it be deemed necessary to facilitate prosecution of the application.

Respectfully submitted,

BANNER & WITCOFF, LTD.

Dated: December 11, 2002

By:


Robert H. Resis

Reg. No. 32,168

Direct Dial: (312) 715-3405


Banner & Witcoff, Ltd.
Ten South Wacker Drive
Chicago, Illinois 60606
(312) 715-1000

VERSION WITH MARKINGS TO SHOW CHANGES MADE

In the Specification:

Please replace the paragraph beginning on page 2, line 11 with the following rewritten paragraph:

--In one form of the compound of Formula (1), the functionalised linkage group Z is selected from the group of halogen or other leaving group, nitro, nitroso, imide, dione of the formula,



vinyl group of formula $\text{Het}^1\text{-C}(\text{Het}^2)=\text{CH}_2$ (where Het^1 and Het^2 are the same or different and is each a nitrogen containing heterocyclic group or Het^1 is a nitrogen containing heterocyclic group and Het^2 is H), $-\text{C}(=\text{NH})\text{OR}^2$, NCO , NCS , COR'' , COOR' , SR^2 , NHN^2R^3 , $\text{NHCONHN}^2\text{R}^3$, $\text{NHCSNHN}^2\text{R}^3$, $[\text{CONR}^2]$ CONR^2R^3 , OR^2 , NR^2R^3 , $(\text{CH}_2)_p\text{R}^1$, $(\text{CH}_2)_p\text{ArR}^1$, $(\text{CH}^2\text{O})_p\text{CH}_2\text{R}^1$, $(\text{CH}_2\text{OCH}_2\text{O})_q\text{ArR}^1$, $(\text{CHCH})_r\text{R}^1$, $(\text{CHCH})_r\text{ArR}^1$ where R^2 and R^3 are the same or different and are independently selected from h, $(\text{CH}_2)_p\text{R}^1$, $(\text{CH}_2)_p\text{ArR}^1$, $(\text{CH}_2\text{O})_p\text{CH}_2\text{R}^1$, $(\text{CH}_2\text{OCH}_2\text{O})_q\text{ArR}^1$, $(\text{CHCH})_r\text{R}^1$, $(\text{CHCH})_r\text{ArR}^1$ and where R^1 is selected from SH, OH, NH_2COOH , NCS , $-\text{N}=\text{N}$, or $-\text{C}(=\text{NH})\text{OCH}_3$, COR'' , where R'' is H, halogen, N_3 , alkoxy, Oar, imidyloxy, imidazoyloxy, alkyl, or alkyl substituted with a halogen or other leaving group,

where p is an integer from 1 to 20, more typically 1 to 10, still more typically 1 to 4, even more typically 1 to 2 and yet more typically 1; q is an integer from 1 to 20, more typically 1 to 10, still more typically 1 to 4, even more typically 1 to 2 and yet more typically 1; r is an integer from 1 to 4, more typically 1 or 2, still more typically 1; and Ar is optionally substituted aryl or optionally substituted aralkyl, provided that when one of X and Y is selected from C-NO₂, C-OH, C-Cl, C-CH₃ or C-NH₂ then the other X or Y substituent cannot be selected from C-NO₂, C-OH, C-Cl or C-NH₂. In moieties of formula (CH₂)_pArR¹, one or more methylene groups may also be replaced with O, S, NH or carbonyl, for example C(O)R¹, CH₂C(O)R¹, NHCH₂R¹, NHC(O)R¹, CH₂OR¹, CH₂C(O)NHR¹, and the like.--